

Hopf-Galois systems and Kashiwara algebras

Cyril Grunspan,

Institut de Recherche Mathématique avancée,
Université Louis Pasteur et CNRS,
7 Rue René Descartes,
67084 Strasbourg Cedex, France

February 1, 2008

Abstract

This article is made up with two parts. In the first part, using a recent result of Schauenburg, one generalizes to the case when objects are faithfully flat over the ground ring, the full equivalence between the notions of Hopf-Galois objects and Hopf-Galois systems. In this last description, one gives explicitly an inverse for a Hopf-Galois object T together with its generalized antipode.

In the second part of the article, one shows that the Kashiwara algebras introduced by Kashiwara in his study of crystal bases form Hopf-Galois systems under the coaction of a quantized enveloping algebra of a Kac-Moody algebra. Their classical limits are examples of Sridharan algebras.

Keywords: Hopf-Galois system, Hopf-Galois extension, Kashiwara algebra, Sridharan algebra, torsor.

1 Introduction

In [G], we have introduced the concept of a quantum torsor which turned out to be an intrinsic reformulation of the old notion of Hopf-Galois extension [Sc2], at least in the case when modules are vector spaces over a field. A quantum torsor is an algebra T equipped with a triple coproduct μ satisfying some natural relations of co-associativity and co-unity. The law μ “encodes” the two structures of bi-algebras of the Hopf algebras co-acting on T . The antipodes are then “encoded” themselves in a torsor endomorphism θ on T whose existence is a consequence of torsor axioms satisfied by μ alone [Sc3]. In this last article, Schauenburg also proved the full equivalence between the two notions of non-commutative torsors and Hopf-Galois objects to the case when modules are faithfully flat over the ground ring.

Independently, Bichon introduced in [B] the concept of Hopf-Galois systems which is based on the idea that a torsor is a groupoid (A, B, T, Z) with two

units. Originally, this idea can be found in [Sc1]. With the help of the Tannaka duality, he proved that if the ground ring is a field, then the two concepts of Hopf-Galois systems and Hopf-Galois extensions are equivalent. However, an inverse Z of T is not given explicitly, nor the “generalized antipode” from Z to T which is the main feature of his set of axioms. Note also that his set of axioms is not symmetric. The advantage is that it is easy to handle with into practice.

In the first part of the article, we show that if T is a (A, B) -bi-Galois object, then the algebra $Z = T^{-1}$ introduced by Schauenburg in [Sc1] can be taken so as to form a Hopf-Galois system (A, B, T, Z) . The result is proved for any faithfully flat Hopf-Galois object T . This generalizes Bichon’s theorem. We give explicitly a formula for the “generalized antipode” in terms of the torsor endomorphism θ of T . Moreover, the result one obtains is symmetric in T and Z . Note that if T is autonomous i.e., if the torsor map θ associated to T is bijective, then one can take $Z = T^{\text{op}}$ as proved in [G].

In the second part of this article, we show that the Kashiwara algebras introduced by Kashiwara in his study of crystal basis [K] form Hopf-Galois systems. The Hopf algebras involved in these systems are standard quantum groups $U_q(\mathfrak{g})$ associated to any Kac-Moody algebra $\mathfrak{g}$: Kashiwara algebras are $U_q(\mathfrak{g})$ -Hopf-Galois objects. We give the explicit torsor laws μ and θ . We show that their classical limits are examples of Sridharan algebras [Sr].

Kashiwara algebras provide examples of (A, B) -bi-Galois objects with two *non-isomorphic* Hopf-algebras A and B and with a *non-trivial* torsor map θ , such that at the classical level, the Hopf algebras associated to the corresponding Hopf-Galois systems are *isomorphic* and such that the torsor maps are *trivial*. Proofs of Sections 3, 4 and 5 are very easy and can be omitted.

2 Faithfully flat Hopf-Galois systems

One recalls Bichon’s axioms of a Hopf-Galois system [B]. Fix k a commutative ring.

Definition 1 *A Hopf-Galois system consists of four non-zero k -module-algebras (A, B, T, Z) with the following axioms.*

- i. *The algebras A and B are bi-algebras.*
- ii. *The algebra T is a (A, B) -bicomodule algebra.*
- iii. *There are algebra morphisms $\gamma : A \longrightarrow T \otimes Z$ and $\delta : B \longrightarrow Z \otimes T$ such that the following diagrams commute:*

$$\begin{array}{ccccccc}
 T & \xrightarrow{\alpha_T} & A \otimes T & A & \xrightarrow{\Delta_A} & A \otimes A & B & \xrightarrow{\Delta_B} & B \otimes B \\
 \beta_T \downarrow & & \downarrow \gamma \otimes T & \gamma \downarrow & & \downarrow A \otimes \gamma & \delta \downarrow & & \downarrow \delta \otimes B \\
 T \otimes B & \xrightarrow{T \otimes \delta} & T \otimes Z \otimes T & T \otimes Z & \xrightarrow{\alpha_T \otimes Z} & A \otimes T \otimes Z & Z \otimes T & \xrightarrow{Z \otimes \beta_T} & Z \otimes T \otimes B
 \end{array}$$

iv. There is a linear map $S_Z : Z \longrightarrow T$ such that the following diagrams commute:

$$\begin{array}{ccc} A & \xrightarrow{\varepsilon_A} k \xrightarrow{u_T} & T \\ \gamma \downarrow & & \uparrow m_T \\ T \otimes Z & \xrightarrow{T \otimes S_Z} & T \otimes T \end{array} \quad \begin{array}{ccc} B & \xrightarrow{\varepsilon_B} k \xrightarrow{u_T} & T \\ \delta \downarrow & & \uparrow m_T \\ Z \otimes T & \xrightarrow{S_Z \otimes T} & T \otimes T \end{array}$$

Note that if (A, B, T, Z) is a Hopf-Galois system, then 1) the bi-algebras A and B are in fact Hopf algebras and 2) the “generalized antipode” $S_Z : Z \longrightarrow T^{\text{op}}$ is then an algebra morphism as proved in [B]. It is easy to show that if (A, B, T, Z) is a Hopf-Galois system, then T is in fact a (A, B) -bi-Galois extension of k . Conversely, using techniques of Tannaka duality, Bichon proved the following theorem.

Theorem 1 *If k is a field and if T is a B -Galois extension of k , then there exists an Hopf algebra A , an algebra Z and a map $S_Z : Z \longrightarrow T$ such that (A, B, T, Z) is a Hopf-Galois system.*

We generalize this theorem to the case when T is a faithfully flat B -Galois extension of k . In fact, we prove more. We prove that any faithfully flat Hopf-Galois objects can be part of a *complete* Hopf-Galois system in the sense we gave in [G] without any hypothesis on θ .

Theorem 2 *Let (T, μ, θ) be a faithfully flat quantum torsor over k . Set $A := H_l(T)$, $B := H_r(T)$ and $\alpha_T : T \longrightarrow A \otimes T$ and $\beta_T : T \longrightarrow T \otimes B$ the two natural morphisms defining a structure of (A, B) -bi-Galois extension of k on T . Recall that A (resp. B) is defined as being a certain sub-algebra of $T \otimes T^{\text{op}}$ (resp. $T^{\text{op}} \otimes T$). Then,*

1. *One has $(B \otimes T)^{\text{co}B} = (T \otimes A)^{\text{co}A}$ as subsets of $T \otimes T \otimes T$.*
2. *There is a natural structure of k -algebra on $Z := (B \otimes T)^{\text{co}B} = (T \otimes A)^{\text{co}A}$ as sub-algebra of $T^{\text{op}} \otimes T \otimes T^{\text{op}}$.*
3. *There is a natural structure of (B, A) -bi-Galois extension of k on Z given by morphisms $\beta_Z := (\Delta_B \otimes T) : Z \longrightarrow B \otimes Z$ and $\alpha_Z := (Z \otimes \Delta_A) : Z \longrightarrow Z \otimes A$.*
4. *If $x \in T$, then $S_T(x) := (\theta \otimes T \otimes \theta) \circ \mu^{\text{op}}(x) \in Z$.*
5. *The map $S_T : T \longrightarrow Z^{\text{op}}$ is an algebra morphism.*
6. *If $x \in Z$, then $(\varepsilon_B \otimes T)(x) = (T \otimes \varepsilon_A)(x)$. We denote by $S_Z(x)$ this common value.*
7. *The map $S_Z : Z \longrightarrow T^{\text{op}}$ is an algebra morphism.*
8. *If $h = x_i \otimes y_i \in B \subset T^{\text{op}} \otimes T$, then $\delta(h) := x_i \otimes y_i^{(1)} \otimes y_i^{(2)} \otimes y_i^{(3)} \in Z \otimes T \subset T^{\text{op}} \otimes T \otimes T^{\text{op}} \otimes T$.*

9. If $h = x_i \otimes y_i \in A \subset T \otimes T^{\text{op}}$, then $\gamma(h) := x_i^{(1)} \otimes x_i^{(2)} \otimes x_i^{(3)} \otimes y_i \in T \otimes Z \subset T \otimes T^{\text{op}} \otimes T \otimes T^{\text{op}}$.

10. The map $\delta : B \longrightarrow Z \otimes T$ is an algebra morphism.

11. The map $\gamma : A \longrightarrow T \otimes Z$ is an algebra morphism.

Moreover, the quadruple (A, B, T, Z) equipped with the morphisms $(a_T, \beta_T, \alpha_Z, \beta_Z, \gamma, \delta, S_T, S_Z)$ is a **complete** Hopf-Galois system in the sense of our definition given in [G] and the quantum torsor associated to this Hopf-Galois system is isomorphic to (T, μ, θ) . In particular, (A, B, T, Z) and (B, A, Z, T) are two Hopf-Galois systems in the sense of Definition 1.

Before starting the proof of Theorem 2, we recall the following facts taken from [G].

a) The Hopf algebra A is the set of $x_i \otimes y_i \in T \otimes T^{\text{op}}$ such that

$$x_i^{(1)} \otimes x_i^{(2)} \otimes \theta(x_i^{(3)}) \otimes y_i = x_i \otimes y_i^{(3)} \otimes y_i^{(2)} \otimes y_i^{(1)}.$$

b) The Hopf algebra B is the set of $x_i \otimes y_i \in T^{\text{op}} \otimes T$ such that

$$x_i \otimes \theta(y_i^{(1)}) \otimes y_i^{(2)} \otimes y_i^{(3)} = x_i^{(3)} \otimes x_i^{(2)} \otimes x_i^{(1)} \otimes y_i.$$

c) The comultiplication on $A \subset T \otimes T^{\text{op}}$ is defined by $\Delta_A(x_i \otimes y_i) = x_i^{(1)} \otimes x_i^{(2)} \otimes x_i^{(3)} \otimes y_i$.

d) The comultiplication on $B \subset T^{\text{op}} \otimes T$ is defined by $\Delta_B(x_i \otimes y_i) = x_i \otimes y_i^{(1)} \otimes y_i^{(2)} \otimes y_i^{(3)}$.

e) The antipode on $A \subset T \otimes T^{\text{op}}$ is defined by $S_A(x_i \otimes y_i) = y_i \otimes \theta(x_i)$.

f) The antipode on $B \subset T^{\text{op}} \otimes T$ is defined by $S_B(x_i \otimes y_i) = \theta(y_i) \otimes x_i$.

g) The counit on $A \subset T \otimes T^{\text{op}}$ is defined by $\eta_T \circ \varepsilon_A(x_i \otimes y_i) = x_i y_i$.

h) The counit on $B \subset T^{\text{op}} \otimes T$ is defined by $\eta_T \circ \varepsilon_B(x_i \otimes y_i) = x_i y_i$.

i) For $x \in T$, $\alpha_T(x) = x^{(1)} \otimes x^{(2)} \otimes x^{(3)} \in A \otimes T \subset T \otimes T^{\text{op}} \otimes T$.

j) For $x \in T$, $\beta_T(x) = x^{(1)} \otimes x^{(2)} \otimes x^{(3)} \in T \otimes B \subset T \otimes T^{\text{op}} \otimes T$.

k) If H is a Hopf algebra and if M and N are two H -right-comodules, then $(M \otimes N)^{\text{co } H}$ is the set of $x_i \otimes y_i \in M \otimes N$ such that $x_i \otimes y_{i,N} \otimes y_{i,H} = x_{i,M} \otimes y_i \otimes S(x_{i,H})$.

l) Similarly, if M and N are two H -left-comodules, then $(M \otimes N)^{\text{co } A}$ is the set of $x_i \otimes y_i \in M \otimes N$ such that $x_{i,H} \otimes x_{i,M} \otimes y_i = S(y_{i,H}) \otimes x_i \otimes y_i$.

Proof In the following, the expression $x \otimes y$ (resp. $x \otimes y \otimes z$) will stand for a sum of terms of the form $x_i \otimes y_j \otimes z_k$.

Let $x \otimes y \otimes z \in (B \otimes T)^{\text{co} B} \subset T \otimes T \otimes T$. By d, f, j and k, we get :

$$x \otimes y \otimes z^{(1)} \otimes z^{(2)} \otimes z^{(3)} = x \otimes y^{(1)} \otimes z \otimes \theta(y^{(3)}) \otimes y^{(2)}. \quad (1)$$

So,

$$x \otimes y^{(1)} \otimes y^{(2)} \otimes \theta(y^{(3)}) \otimes z = x \otimes y \otimes z^{(3)} \otimes z^{(2)} \otimes z^{(1)}.$$

So, by a and the fact that T is faithfully flat, we see that $x_i \otimes y_j \otimes z_k \in T \otimes A$. Moreover, by b, we get:

$$x^{(3)} \otimes x^{(2)} \otimes x^{(1)} \otimes y \otimes z = x \otimes \theta(y^{(1)}) \otimes y^{(2)} \otimes y^{(3)} \otimes z. \quad (2)$$

So,

$$x^{(1)} \otimes x^{(2)} \otimes x^{(3)} \otimes y \otimes z = y^{(2)} \otimes \theta(y^{(1)}) \otimes x \otimes y^{(3)} \otimes z.$$

So, by c, e, i and l, we deduce that $x \otimes y \otimes z \in (T \otimes A)^{\text{co} A}$. Therefore, $(B \otimes T)^{\text{co} B} \subset (T \otimes A)^{\text{co} A}$ and similarly, $(T \otimes A)^{\text{co} A} \subset (B \otimes T)^{\text{co} B}$. So, **1) is proved**. Assertions **2) and 3) are proved in [Sc1]**.

Let $x \in T$. By torsor axioms, we have

$$x^{(1)} \otimes x^{(2)} \otimes \theta(x^{(3)}) \otimes x^{(4)} \otimes x^{(5)} = x^{(1)} \otimes x^{(2)(3)} \otimes x^{(2)(2)} \otimes x^{(2)(1)} \otimes x^{(3)} \quad (3)$$

So,

$$\theta(x^{(5)}) \otimes \theta(x^{(4)}) \otimes \theta(x^{(3)}) \otimes x^{(2)} \otimes \theta(x^{(1)}) = \theta(x^{(3)}) \otimes \theta(x^{(2)})^{(1)} \otimes x^{(2)(2)} \otimes x^{(2)(3)} \otimes \theta(x^{(1)}).$$

So,

$$(\mu^{\text{op}} \otimes T \otimes T)(\theta(x^{(3)}) \otimes x^{(2)} \otimes \theta(x^{(1)})) = (T \otimes \theta \otimes T \otimes T \otimes T) \circ (T \otimes \mu \otimes T)(\theta(x^{(3)}) \otimes x^{(2)} \otimes \theta(x^{(1)}))$$

Hence, by b, we see that $\theta(x^{(3)}) \otimes x^{(2)} \otimes \theta(x^{(1)}) \in B \otimes T$. On the other hand, by (3), we get

$$\theta(x^{(5)}) \otimes x^{(4)} \otimes \theta(x^{(1)}) \otimes \theta(x^{(2)}) \otimes \theta(x^{(3)}) = \theta(x^{(3)}) \otimes x^{(2)(1)} \otimes \theta(x^{(1)}) \otimes \theta(x^{(2)(3)}) \otimes x^{(2)(2)}.$$

So, by d, f, j and k, $\theta(x^{(3)}) \otimes x^{(2)} \otimes \theta(x^{(1)}) \in (B \otimes T)^{\text{co} B}$ and **4) is proved**.

For $x, y \in T$, we have $S_T(y) \cdot S_T(x) = \theta(x^{(3)})\theta(y^{(3)}) \otimes y^{(2)}x^{(2)} \otimes \theta(x^{(1)}) \otimes \theta(y^{(1)})$ since $B \otimes T$ is seen as a subalgebra of $T^{\text{op}} \otimes T \otimes T^{\text{op}}$. Thus, $S_T(y) \cdot S_T(x) = \theta(x^{(3)}y^{(3)}) \otimes y^{(2)}x^{(2)} \otimes \theta(x^{(1)}y^{(1)}) = S_T(x \cdot y)$. So, $S_T : T \longrightarrow Z^{\text{op}}$ is an algebra morphism and **5) is proved**.

Let $u =: x_i \otimes y_j \otimes z_k \in Z$. Then, by g and h, $(\varepsilon_B \otimes T)(u) = x_i y_j z_k = (T \otimes \varepsilon_A)(u)$. So, **6) is proved**.

Suppose that $u =: x \otimes y \otimes z \in Z$ and $v =: x' \otimes y' \otimes z' \in Z$. Then, by h and 2),

$$\begin{aligned} 1 \otimes S_Z(v) S_Z(u) &= 1 \otimes \varepsilon_B(x' \otimes y') z' \cdot \varepsilon_B(x \otimes y) z = x' y' \otimes z' \cdot \varepsilon_B(x \otimes y) z \\ &= x' \varepsilon_B(x \otimes y) y' \otimes z' z = x' x y y' \otimes z' z \\ &= 1 \otimes S_Z(x' x \otimes y y' \otimes z' z) = 1 \otimes S_Z(uv) \end{aligned}$$

Hence, $S_Z(uv) = S_Z(vu)$ and **7) is proved**.

Let $h = x_i \otimes y_i \in B \subset T^{\text{op}} \otimes T$. By, d we already have $\delta(h) := x \otimes y^{(1)} \otimes y^{(2)} \otimes y^{(3)} \in B \otimes B \subset B \otimes T \otimes T$. Moreover, $\delta(h) \in (B \otimes T)^{\text{co}B} \otimes T$ since by torsor axioms, we have

$$x \otimes y^{(1)} \otimes y^{(2)(1)} \otimes y^{(2)(2)} \otimes y^{(2)(3)} \otimes y^{(3)} = x \otimes y^{(1)} \otimes y^{(4)} \otimes \theta(x^{(3)}) \otimes y^{(2)} \otimes y^{(5)}.$$

So, **8) is proved**. In the same way, **9) is true**.

From the definition of δ and γ , it is easy to see that they define algebra morphisms. So, **10) and 11) are true**.

Let us prove that (A, B, T, Z) is a Hopf-Galois system in the sense of Definition 1. Axioms (i) and (ii) of Definition 1 are clearly satisfied. Let $x \in T$. Then,

$$\begin{aligned} (\gamma \otimes T) \circ \alpha_T(x) &= (\gamma \otimes T)(x^{(1)} \otimes x^{(2)} \otimes x^{(3)}) \\ &= x^{(1)} \otimes x^{(2)} \otimes x^{(3)} \otimes x^{(4)} \otimes x^{(5)} \\ &= (T \otimes \delta)(x^{(1)} \otimes x^{(2)} \otimes x^{(3)}) \\ &= (T \otimes \delta) \circ \beta_T(x). \end{aligned}$$

This proves the commutativity of the first diagram of (iii) in Definition 1. Let $x \otimes y \in A$. Then,

$$\begin{aligned} (A \otimes \gamma) \circ \Delta_A(x \otimes y) &= (A \otimes \gamma)(x^{(1)} \otimes x^{(2)} \otimes x^{(3)} \otimes y) \\ &= x^{(1)} \otimes x^{(2)} \otimes x^{(3)} \otimes x^{(4)} \otimes x^{(5)} \otimes y \\ &= (\alpha_T \otimes Z)(x^{(1)} \otimes x^{(2)} \otimes x^{(3)} \otimes y) \\ &= (\alpha_T \otimes Z) \circ \gamma(x \otimes y) \end{aligned}$$

and

$$\begin{aligned} m_T \circ (T \otimes S_Z) \circ \gamma(x \otimes y) &= m_T \circ (T \otimes S_Z) \circ \gamma(x^{(1)} \otimes x^{(2)} \otimes x^{(3)} \otimes y) \\ &= m_T \circ (x^{(1)} \otimes \varepsilon_B(x^{(2)} \otimes x^{(3)})y) \\ &= m_T(x \otimes y) \\ &= \eta_T \circ \varepsilon_A(x \otimes y) \end{aligned}$$

Moreover, for $x \otimes y \in B$, we have,

$$\begin{aligned} (\delta \otimes B) \circ \Delta_B(x \otimes y) &= (\delta \otimes B)(x \otimes y^{(1)} \otimes y^{(2)} \otimes y^{(3)}) \\ &= x \otimes y^{(1)} \otimes y^{(2)} \otimes y^{(3)} \otimes y^{(4)} \otimes y^{(5)} \\ &= (Z \otimes \beta_T)(x \otimes y^{(1)} \otimes y^{(2)} \otimes y^{(3)}) \\ &= (Z \otimes \beta_T) \circ \delta(x \otimes y) \end{aligned}$$

and

$$\begin{aligned} m_T \circ (S_Z \otimes T) \circ \delta(x \otimes y) &= m_T \circ (S_Z \otimes T)(x \otimes y^{(1)} \otimes y^{(2)} \otimes y^{(3)}) \\ &= m_T(x \varepsilon_A(y^{(1)} \otimes y^{(2)}) \otimes y^{(3)}) \\ &= m_T(x \otimes y) \\ &= \eta_T \circ e_B(x \otimes y). \end{aligned}$$

All these identities above show that (A, B, T, Z) is a **Hopf-Galois system**.

Let us prove that (B, A, Z, T) equipped with the maps $(\alpha_Z, \beta_Z, \dots, S_T)$ is also a Hopf-Galois system. By 3), Axioms (i) and (ii) for this system are satisfied. Let $x \otimes y \otimes z \in Z$. Then,

$$\begin{aligned}
(Z \otimes \gamma) \circ \alpha_Z(x \otimes y \otimes z) &= (Z \otimes \gamma)(x \otimes y^{(1)} \otimes y^{(2)} \otimes y^{(3)} \otimes z) \\
&= x \otimes y^{(1)} \otimes y^{(2)} \otimes y^{(3)(1)} \otimes y^{(3)(2)} \otimes y^{(3)(3)} \otimes z \\
&= x \otimes y^{(1)(1)} \otimes y^{(1)(2)} \otimes y^{(1)(3)} \otimes y^{(2)} \otimes y^{(3)} \otimes z \\
&= (\delta \otimes Z) \circ \beta_Z(x \otimes y \otimes z)
\end{aligned}$$

Moreover, for $x \otimes y \in A$, we have

$$\begin{aligned}
(\gamma \otimes A) \circ \Delta_A(x \otimes y) &= (\gamma \otimes A)(x^{(1)} \otimes x^{(2)} \otimes x^{(3)} \otimes y) \\
&= x^{(1)(1)} \otimes x^{(1)(2)} \otimes x^{(1)(3)} \otimes x^{(2)} \otimes x^{(3)} \otimes y \\
&= x^{(1)} \otimes x^{(2)} \otimes x^{(3)(1)} \otimes x^{(3)(2)} \otimes x^{(3)(3)} \otimes y \\
&= (T \otimes \alpha_Z)(x^{(1)} \otimes x^{(2)} \otimes x^{(3)} \otimes y) \\
&= (T \otimes A) \circ (T \otimes \alpha_Z)(x \otimes y)
\end{aligned}$$

and

$$\begin{aligned}
m_Z \circ (S_T \otimes Z) \circ \gamma(x \otimes y) &= m_Z \circ (S_T \otimes Z)(x^{(1)} \otimes x^{(2)} \otimes x^{(3)} \otimes y) \\
&= m_Z \circ (\theta(x^{(3)}) \otimes x^{(2)} \otimes \theta(x^{(1)}) \otimes x^{(4)} \otimes x^{(5)} \otimes y) \\
&= x^{(4)} \theta(x^{(3)}) \otimes x^{(2)} x^{(5)} \otimes y \theta(x^{(1)}) \\
&= 1 \otimes x^{(2)} x^{(3)} \otimes y \theta(x^{(1)}) \\
&= 1 \otimes 1 \otimes y \theta(x) \\
&= \eta_Z \circ \varepsilon_A(x \otimes y)
\end{aligned}$$

We also see that for $x \otimes y \in B$, we have

$$\begin{aligned}
(B \otimes \delta) \circ \Delta_B(x \otimes y) &= (B \otimes \delta)(x \otimes y^{(1)} \otimes y^{(2)} \otimes y^{(3)}) \\
&= x \otimes y^{(1)} \otimes y^{(2)} \otimes y^{(3)} \otimes y^{(4)} \otimes y^{(5)} \\
&= (\beta_Z \otimes T)(x \otimes y^{(1)} \otimes y^{(2)} \otimes y^{(3)}) \\
&= (\beta_Z \otimes T) \circ \delta(x \otimes y)
\end{aligned}$$

and

$$\begin{aligned}
m_Z \circ (Z \otimes S_T) \circ \delta(x \otimes y) &= m_Z \circ (Z \otimes S_T)(x \otimes y^{(1)} \otimes y^{(2)} \otimes y^{(3)}) \\
&= m_Z(x \otimes y^{(1)} \otimes y^{(2)} \otimes \theta(y^{(5)}) \otimes y^{(4)} \otimes \theta(y^{(3)})) \\
&= \theta(y^{(5)}) x \otimes y^{(1)} y^{(4)} \otimes \theta(y^{(3)}) y^{(2)} \\
&= \theta(y^{(3)}) x \otimes y^{(1)} y^{(2)} \otimes 1 \\
&= \theta(y) x \otimes 1 \otimes 1 \\
&= \eta_Z \circ \varepsilon_B(x \otimes y)
\end{aligned}$$

We then deduce that (B, A, Z, T) is a **Hopf-Galois system**. To end the proof, we need to check some compatibility relations between the Hopf-Galois systems (A, B, T, Z) and (B, A, Z, T) . Let $x \otimes y \in A$. Then,

$$\begin{aligned}
(\beta_T \otimes Z) \circ \gamma(x \otimes y) &= (\beta_T \otimes Z)(x^{(1)} \otimes x^{(2)} \otimes x^{(3)} \otimes y) \\
&= x^{(1)} \otimes x^{(2)} \otimes x^{(3)} \otimes x^{(4)} \otimes x^{(5)} \otimes y \\
&= (T \otimes \beta_Z)(x^{(1)} \otimes x^{(2)} \otimes x^{(3)} \otimes y) \\
&= (T \otimes \beta_Z) \circ \gamma(x \otimes y)
\end{aligned}$$

and

$$\begin{aligned}
\tau_{(T,Z)} \circ (S_T \otimes S_Z) \circ \gamma(x \otimes y) &= \tau_{(T,Z)} \circ (S_T \otimes S_Z)(x^{(1)} \otimes x^{(2)} \otimes x^{(3)} \otimes y) \\
&= \tau_{(T,Z)}(\theta(x^{(3)}) \otimes x^{(2)} \otimes \theta(x^{(1)}) \otimes \varepsilon_B(x^{(4)} \otimes x^{(5)})y) \\
&= \tau_{(T,Z)}(\theta(x^{(3)}) \otimes x^{(2)} \otimes \theta(x^{(1)}) \otimes y) \\
&= y \otimes \theta(x^{(3)}) \otimes x^{(2)} \otimes \theta(x^{(1)}) \\
&\stackrel{a}{=} y^{(1)} \otimes y^{(2)} \otimes y^{(3)} \otimes \theta(x) \\
&= \gamma(y \otimes \theta(x)) = \gamma \circ S_A(x \otimes y)
\end{aligned}$$

Similarly, we prove $(\alpha_Z \otimes T) \circ \delta = (Z \otimes \alpha_T) \circ \delta$ and $\tau_{(Z,T)} \circ (S_Z \otimes S_T) \circ \delta = \delta \circ S_B$. Let $x \in T$. Then,

$$\begin{aligned}
\tau_{(A,Z)} \circ (S_A \otimes S_T) \circ \alpha_T(x) &= \tau_{(A,Z)} \circ (S_A \otimes S_T)(x^{(1)} \otimes x^{(2)} \otimes x^{(3)}) \\
&= \tau_{(A,Z)}(x^{(2)} \otimes \theta(x^{(1)}) \otimes \theta(x^{(5)}) \otimes x^{(4)} \otimes \theta(x^{(3)})) \\
&= \theta(x^{(5)}) \otimes x^{(4)} \otimes \theta(x^{(3)}) \otimes x^{(2)} \otimes \theta(x^{(1)}) \\
&= \theta(x^{(3)}) \otimes x^{(2)(1)} \otimes x^{(2)(2)} \otimes x^{(2)(3)} \otimes \theta(x^{(1)}) \\
&= \alpha_Z(\theta(x^{(3)}) \otimes x^{(2)} \otimes \theta(x^{(1)})) \\
&= \alpha_Z \circ S_T(x).
\end{aligned}$$

In the same way, we prove that $\tau_{(Z,B)} \circ (S_T \otimes S_B) \circ \beta_T = \beta_Z \circ S_T$. Moreover, for $x \otimes y \otimes z \in Z$, we have

$$\begin{aligned}
\beta_T \circ S_Z(x \otimes y \otimes z) &= \beta_T(x \otimes \varepsilon_A(y \otimes z)) \\
&= x^{(1)} \otimes x^{(2)} \otimes x^{(3)} \otimes \varepsilon_A(y \otimes z) \\
&= y^{(2)} \otimes \theta(y^{(1)}) \otimes x \otimes \varepsilon_A(y^{(3)} \otimes z) \\
&= y^{(2)} \otimes \varepsilon_A(y^{(3)} \otimes z) \otimes \theta(y^{(1)}) \otimes x \\
&= \tau_{(B,T)} \circ (S_B \otimes S_Z)(x \otimes y^{(1)} \otimes y^{(2)} \otimes y^{(3)} \otimes z) \\
&= \tau_{(B,T)} \circ (S_B \otimes S_Z) \circ \beta_Z(x \otimes y \otimes z)
\end{aligned}$$

In the same way, we prove that $\alpha_T \circ S_Z = \tau_{(T,A)} \circ (S_Z \otimes S_A) \circ \alpha_Z$. Therefore, all the axioms of compatibility between the two Hopf-Galois systems (A, B, T, Z) and (B, A, Z, T) are satisfied (see [G]). Thus, (A, B, T, Z) is a **complete Hopf-Galois system**.

□

Given the full equivalence between Hopf-Galois extensions and quantum torsors as proved in [Sc3], we can state the following generalization of Bichon's theorem.

Corollary 1 *If T is a faithfully flat B -Galois extension of k , then there exists an Hopf algebra A , an algebra Z and algebra morphisms $S_T : T \rightarrow Z^{\text{op}}$ and $S_Z : Z \rightarrow T^{\text{op}}$ such that (A, B, T, Z) is a complete Hopf-Galois system in the sense we gave in [G]. In particular, (A, B, T, Z) and (B, A, Z, T) are two Hopf-Galois systems in the sense of Definition 1.*

3 Sridharan algebras and $\text{Bigal}(U(\mathfrak{g}))$

In this short section, we introduce the Sridharan algebras [Sr] and we prove that they are $U(\mathfrak{g})$ -Galois objects.

Definition 2 *Let $(\mathfrak{g}, [,])$ be a $\mathbb{Q}$ -Lie algebra and let c be a 2-cocycle of $\mathfrak{g}$ with values in $\mathbb{Q}$ i.e., c satisfies the following equation:*

$$c([x, y], z) + c([y, z], x) + c([z, x], y) = 0$$

for all $x, y, z \in \mathfrak{g}$. The Sridharan algebra $U_c(\mathfrak{g})$ associated to c is $T(\mathfrak{g})/I_c(\mathfrak{g})$ where $T(\mathfrak{g})$ is the tensorial algebra of $\mathfrak{g}$ and $I_c(\mathfrak{g})$ is the ideal generated by elements $x \cdot y - y \cdot x - [x, y] - c(x, y) \cdot 1$ with $x, y \in \mathfrak{g}$.

It can be shown that two cocycles define two isomorphic Sridharan algebras if and only if the two cocycles are cohomologous. The point is that Sridharan are examples of $U(\mathfrak{g})$ -torsors.

Theorem 3 *Let $\mathfrak{g}$ be a $\mathbb{Q}$ -Lie algebra and c a 2-cocycle of $\mathfrak{g}$ with values in $\mathbb{Q}$. Then, $(U_c(\mathfrak{g}), U_{-c}(\mathfrak{g}))$ is a $(U(\mathfrak{g}), U(\mathfrak{g}))$ -Hopf-Galois system over $\mathbb{Q}$. The maps defining this Hopf-Galois system are all of the form $x \mapsto 1 \otimes x + x \otimes 1$ for $x \in \mathfrak{g}$ and $S(x) = -x, x \in \mathfrak{g}$ for the generalized antipodes.*

By [G], we then deduce the following corollaries.

Corollary 2 *Let $\mathfrak{g}$ be a $\mathbb{Q}$ -Lie algebra and c a 2-cocycle of $\mathfrak{g}$ with values in $\mathbb{Q}$. The Sridharan algebra $U_c(\mathfrak{g})$ is a quantum torsor and $H_l(U_c(\mathfrak{g})) \cong H_r(U_c(\mathfrak{g})) \cong U(\mathfrak{g})$. The torsor laws are defined by $\theta_{U_c(\mathfrak{g})} = \text{Id}_{U_c(\mathfrak{g})}$ and $\mu(x) = x \otimes 1 \otimes 1 - 1 \otimes x \otimes 1 + 1 \otimes 1 \otimes x$ for $x \in \mathfrak{g}$.*

Corollary 3 *Sridharan algebras are $(U(\mathfrak{g}), U(\mathfrak{g}))$ -Hopf-bi-Galois extensions of $\mathbb{Q}$.*

Proposition 1 *Let $\mathfrak{g}$ be a $\mathbb{Q}$ -Lie algebra and c, c' two 2-cocycles of $\mathfrak{g}$ with values in $\mathbb{Q}$. Then, $U_c(\mathfrak{g}) \square_{U(\mathfrak{g})} U_{c'}(\mathfrak{g}) \cong U_{c \cdot c'}(\mathfrak{g})$, where $A \square_H B$ denotes the H -coproduct of a H -right-comodule-algebra A and a H -left-comodule-algebra B . The isomorphism is an isomorphism of $(U(\mathfrak{g}), U(\mathfrak{g}))$ -Hopf-bi-Galois extensions of $\mathbb{Q}$.*

Corollary 4 *The map $c \longrightarrow U_c(\mathfrak{g})$ defines a monomorphism of groups of $H^2(\mathfrak{g})$ into $\text{Bigal}(U(\mathfrak{g}))$.*

We conjecture that $\text{Bigal}(U(\mathfrak{g})) = H^2(\mathfrak{g})$. This result is may be known of specialists but we are unable to give a reference.

4 Kashiwara algebras

In Section 4.2, we recall the definition of Kashiwara algebras and we prove that they can be equipped with a quantum torsor structure. Then, in Section 4.4, we will identify the two Hopf algebras associated with these quantum torsors. First, we fix some notations.

4.1 Notations.

We fix $\mathfrak{g}$ a symmetrizable Kac-Moody algebra over $\mathbb{Q}$ with a Cartan sub-algebra $\mathfrak{t}$, $\{\alpha_i \in \mathfrak{t}^*\}_{i \in I}$ the set of simple roots and $\{h_i \in \mathfrak{t}\}_{i \in I}$ the set of coroots, where I is a finite index set. We define an inner product on $\mathfrak{t}^*$ such that $(\alpha_i, \alpha_i) \in \mathbb{N}$ and $\langle h_i, \lambda \rangle = 2(\alpha_i, \lambda)/(\alpha_i, \alpha_i)$ for $\lambda \in \mathfrak{t}^*$. Set $Q = \oplus_i \mathbb{Z}\alpha_i$, $P = \oplus_i \mathbb{Z}h_i$ and $P^* = \{h \in \mathfrak{t}/\langle h, P \rangle \subset \mathbb{Z}\}$. We set $q = \exp(\hbar)$, $q_i = q^{\langle \alpha_i, \alpha_i \rangle / 2}$, $t_i = q^{h_i}$, $[n]_i = (q_i^n - q_i^{-n})/(q_i - q_i^{-1})$, $[n]_i! = \prod_{k=1}^n [k]_i$. The ground ring R is $\mathbb{Q}[[\hbar]]$. We denote by F the field of Laurent series $\mathbb{Q}((\hbar))$. If M is a R -module, we will set $M_F := M \otimes_R F$.

4.2 Kashiwara algebras and quantum torsors

Usually, Kashiwara algebras are defined over F . However, it can be useful to see them as topologically free R -modules, especially because we will examine their classical limits.

Definition 3 *The Kashiwara algebra $B_q(\mathfrak{g})$ [?] is the associative R -algebra generated by generators $e'_i, f_i, i \in I, q^h, h \in P$ and relations :*

$$q^h e'_i q^{-h} = q^{\langle h, \alpha_i \rangle} e'_i \quad (4)$$

$$q^h f_i q^{-h} = q^{-\langle h, \alpha_i \rangle} f_i \quad (5)$$

$$e'_i f_j = q_i^{\langle h_i, \alpha_j \rangle} f_j e'_i + \delta_{i,j}. \quad (6)$$

$$\sum_{k=0}^{1-\langle h_i, \alpha_j \rangle} (-1)^k X_i^{(k)} X_j X_i^{(1-\langle h_i, \alpha_j \rangle)} = 0$$

with $X = e', f$ and $X_i^{(n)} = X_i^n / [n]_i!$. We denote in brief B for $B_q(\mathfrak{g})$.

Proposition 2 *The algebra B is a topologically free R -module. A basis for B over R is given by the family $(e'_{i_1})^\alpha q^h f_{i_2}^\beta$ with $\alpha, \beta \in \mathbb{N}$, $i_1, i_2 \in I$ and $h \in P$.*

The following theorem proves the existence of a quantum torsor structure on B .

Theorem 4 *The morphisms μ_B and θ_B below are well defined and (B, μ_B, θ_B) is an autonomous quantum torsor $[G]$.*

$$\begin{aligned} \mu_B : \quad B &\longrightarrow B \otimes B^{\text{op}} \otimes B \\ e'_i &\longmapsto 1 \otimes 1 \otimes e'_i - 1 \otimes e'_i t_i \otimes t_i^{-1} + e'_i \otimes t_i \otimes t_i^{-1} \\ f_i &\longmapsto 1 \otimes 1 \otimes f_i - 1 \otimes f_i t_i \otimes t_i^{-1} + f_i \otimes t_i \otimes t_i^{-1} \\ q^h &\longmapsto q^h \otimes q^{-h} \otimes q^h \end{aligned}$$

$$\begin{aligned} \theta_B : \quad B &\longrightarrow B \\ e'_i &\longmapsto t_i^{-1} e'_i t_i \\ f_i &\longmapsto t_i^{-1} f_i t_i \\ q^h &\longmapsto q^h \end{aligned}$$

We want now to identify the two Hopf algebras $H_l(B)$ and $H_r(B)$ associated with the quantum torsor (B, μ_B, θ_B) . To this end, we need to define some quantum groups.

4.3 The quantum groups $U_q(\mathfrak{g})$, $U'_q(\mathfrak{g})$ and $\hat{U}_q(\mathfrak{g})$

In this section, we will define three quantum groups which will turn out to coact on B . The first one is the Drinfeld-Jimbo standard quantum group.

Definition 4 *The quantum group $U := U_q(\mathfrak{g})$ is the associative R -algebra over R generated by e_i, f_i, q^h ($i \in I, h \in P^*$) submitted to the following relations:*

$$q^h e_i q^{-h} = q^{\langle h, \alpha_i \rangle} e_i \quad (7)$$

$$q^h f_i q^{-h} = q^{-\langle h, \alpha_i \rangle} f_i \quad (8)$$

$$[e_i, f_j] = \delta_{i,j} (t_i - t_i^{-1}) / (q_i - q_i^{-1}) \quad (9)$$

$$\sum_{k=0}^{1-\langle h_i, \alpha_j \rangle} (-1)^k X_i^{(k)} X_j X_i^{(1-\langle h_i, \alpha_j \rangle)} = 0 \quad (10)$$

with $X = e, f$ and $X_i^{(n)} = X_i^n / [n]_i!$.

The co-multiplication Δ on U is the algebra homomorphism defined by:

$$\Delta(q^h) = q^h \otimes q^h \quad (11)$$

$$\Delta(e_i) = e_i \otimes 1 + t_i \otimes e_i \quad (12)$$

$$\Delta(f_i) = f_i \otimes t_i^{-1} + 1 \otimes f_i. \quad (13)$$

The antipode $S : U \longrightarrow U$ is the anti-isomorphism defined by the formulas:

$$S(e_i) = -t_i^{-1} e_i \quad (14)$$

$$S(f_i) = -f_i \quad (15)$$

$$S(q^h) = q^{-h}. \quad (16)$$

The counit $\varepsilon : U \longrightarrow R$ is the algebra homomorphism defined by:

$$\varepsilon(\mathbf{e}_i) = 0 \quad (17)$$

$$\varepsilon(\mathbf{f}_i) = 0 \quad (18)$$

$$\varepsilon(\mathbf{q}^h) = 1. \quad (19)$$

We define now the two quantum groups $U'_q(\mathfrak{g})$ and $\hat{U}_q(\mathfrak{g})$. They correspond to two different quantizations of a common enveloping algebra $U(\mathfrak{a})$ where $\mathfrak{a}$ is the Lie algebra defined below.

Definition 5 Let $\mathfrak{b}_+$ and $\mathfrak{b}_-$ be the two Borel parts of $\mathfrak{g}$. We denote by $\mathfrak{a}$ the sub-Lie algebra of pairs $(x_+, x_-) \in \mathfrak{b}_+ \times \mathfrak{b}_-$ such that there is $h \in \mathfrak{t}$ satisfying $x_{\pm} - h \in \mathfrak{n}_{\pm}$.

Definition 6 The quantum group $\hat{U}_q(\mathfrak{g})$ (denoted in brief by $\hat{U}$) is the associative algebra generated by generators $\hat{\mathbf{e}}_i, \mathbf{f}_i, i \in I$ and $\mathbf{q}^h, h \in P^*$ satisfying

$$\mathbf{q}^h \hat{\mathbf{e}}_i \mathbf{q}^{-h} = \mathbf{q}^{\langle h, \alpha_i \rangle} \hat{\mathbf{e}}_i \quad (20)$$

$$\mathbf{q}^h \mathbf{f}_i \mathbf{q}^{-h} = \mathbf{q}^{-\langle h, \alpha_i \rangle} \mathbf{f}_i \quad (21)$$

$$[\hat{\mathbf{e}}_i, \mathbf{f}_j] = \delta_{i,j} (\mathbf{t}_i - \mathbf{t}_i^{-1}) \quad (22)$$

$$\sum_{k=0}^{1-\langle h_i, \alpha_j \rangle} (-1)^k \mathbf{X}_i^{(k)} \mathbf{X}_j \mathbf{X}_i^{(1-\langle h_i, \alpha_j \rangle)} = 0 \quad (23)$$

with $\mathbf{X} = \mathbf{e}, \mathbf{f}$ and $\mathbf{X}_i^{(n)} = \mathbf{X}_i^n / [n]_i!$.

The co-multiplication $\hat{\Delta}$ on $\hat{U}$ is the algebra homomorphism defined by :

$$\hat{\Delta}(\mathbf{q}^h) = \mathbf{q}^h \otimes \mathbf{q}^h \quad (24)$$

$$\hat{\Delta}(\hat{\mathbf{e}}_i) = \hat{\mathbf{e}}_i \otimes 1 + \mathbf{t}_i \otimes \hat{\mathbf{e}}_i \quad (25)$$

$$\hat{\Delta}(\mathbf{f}_i) = \mathbf{f}_i \otimes \mathbf{t}_i^{-1} + 1 \otimes \mathbf{f}_i. \quad (26)$$

The antipode $\hat{S} : \hat{U} \longrightarrow \hat{U}$ is the anti-isomorphism defined by the formulas:

$$\hat{S}(\hat{\mathbf{e}}_i) = -\mathbf{t}_i^{-1} \hat{\mathbf{e}}_i \quad (27)$$

$$\hat{S}(\mathbf{f}_i) = -\mathbf{f}_i \quad (28)$$

$$\hat{S}(\mathbf{q}^h) = \mathbf{q}^{-h}. \quad (29)$$

The counit $\hat{\varepsilon} : \hat{U} \longrightarrow R$ is the algebra homomorphism defined by:

$$\hat{\varepsilon}(\hat{\mathbf{e}}_i) = 0 \quad (30)$$

$$\hat{\varepsilon}(\mathbf{f}_i) = 0 \quad (31)$$

$$\hat{\varepsilon}(\mathbf{q}^h) = 1. \quad (32)$$

It can be shown that the above relations defines effectively a Hopf algebra. *A priori*, the common notation $\mathbf{f}_i$ for the generators of U and $\hat{U}$ could be ambiguous for further computations. It is not the case as expressed by the following proposition which links $\hat{U}$ with U .

Proposition 3 *The morphism from $\hat{U}$ to U which sends $\hat{e}_i, \mathbf{f}_i, \mathbf{q}^h$ to $(q_i - q_i^{-1})e_i, \mathbf{f}_i, \mathbf{q}^h$ is a monomorphism of Hopf-algebras.*

From this, we deduce easily the following corollary.

Corollary 5 *The quantum group $\hat{U}$ is a F -form of $U : \hat{U} \otimes F = U \otimes F$*

Next, we turn to $U'_q(\mathfrak{g})$.

Definition 7 *The quantum group $U'_q(\mathfrak{g})$ (denoted in brief by U') is the associative R -algebra over R generated by generators $e'_i, \mathbf{f}'_i, i \in I$ and $\mathbf{q}^h, h \in P^*$ submitted to the following relations:*

$$\begin{aligned} \mathbf{q}^h e'_i \mathbf{q}^{-h} &= \mathbf{q}^{\langle h, \alpha_i \rangle} e'_i \\ \mathbf{q}^h \mathbf{f}'_i \mathbf{q}^{-h} &= \mathbf{q}^{-\langle h, \alpha_i \rangle} \mathbf{f}'_i \\ e'_i \mathbf{f}'_j &= \mathbf{q}_i^{-\langle h_i, \alpha_j \rangle} \mathbf{f}'_j e'_i \end{aligned}$$

$$\sum_{k=0}^{1-\langle h_i, \alpha_j \rangle} (-1)^k \mathbf{X}_i^{(k)} \mathbf{X}_j \mathbf{X}_i^{(1-\langle h_i, \alpha_j \rangle)} = 0$$

with $\mathbf{X} = e', \mathbf{f}'$ and $\mathbf{X}_i^{(n)} = \mathbf{X}_i^n / [n]_i!$. The co-multiplication Δ' on U' is the algebra homomorphism defined by :

$$\Delta'(\mathbf{q}^h) = \mathbf{q}^h \otimes \mathbf{q}^h \quad (33)$$

$$\Delta'(e'_i) = e'_i \otimes 1 + \mathbf{t}_i \otimes e'_i \quad (34)$$

$$\Delta'(\mathbf{f}'_i) = \mathbf{f}'_i \otimes 1 + \mathbf{t}_i \otimes \mathbf{f}'_i. \quad (35)$$

The antipode $S' : U' \longrightarrow U'$ is the anti-isomorphism defined by the formulas:

$$S'(e'_i) = -\mathbf{t}_i^{-1} e'_i \quad (36)$$

$$S'(\mathbf{f}'_i) = \mathbf{t}_i^{-1} \mathbf{f}'_i \quad (37)$$

$$S'(\mathbf{q}^h) = \mathbf{q}^{-h}. \quad (38)$$

The counit $\varepsilon' : U' \longrightarrow R$ is the algebra homomorphism defined by:

$$\varepsilon'(e'_i) = 0 \quad (39)$$

$$\varepsilon'(\mathbf{f}'_i) = 0 \quad (40)$$

$$\varepsilon'(\mathbf{q}^h) = 1. \quad (41)$$

Here also, it can be shown that the above relations defines effectively a Hopf algebra. We are now ready to identify $H_l(B)$ and $H_r(B)$.

4.4 Kashiwara algebras and Hopf-Galois systems

4.4.1 Hopf Galois systems and Kashiwara algebras over R

We are going to embed (B, B^{op}) into a structure of autonomous Hopf-Galois systems.

Proposition 4 *The morphisms $\alpha_B, \beta_B, \alpha_{B^{\text{op}}}, \beta_{B^{\text{op}}}, \gamma, \delta, S_B, S_{B^{\text{op}}}$ below are well defined and $(U', \hat{U}, B, B^{\text{op}})$ is a (complete) autonomous Hopf-Galois system.*

$$\begin{aligned}
\alpha_B : \quad & B \longrightarrow U' \otimes B \\
& e'_i \longmapsto 1 \otimes e'_i + t_i^{-1} e'_i \otimes t_i^{-1} \\
& f_i \longmapsto 1 \otimes f_i + t_i^{-1} f'_i \otimes t_i^{-1} \\
& q^h \longmapsto q^h \otimes q^h \\
\\
\beta_B : \quad & B \longrightarrow B \otimes \hat{U} \\
& e'_i \longmapsto 1 \otimes t_i^{-1} \hat{e}_i + e'_i \otimes t_i^{-1} \\
& f_i \longmapsto f_i \otimes t_i^{-1} + 1 \otimes f_i \\
& q^h \longmapsto q^h \otimes q^h \\
\\
\alpha_{B^{\text{op}}} : \quad & B^{\text{op}} \longrightarrow \hat{U} \otimes B^{\text{op}} \\
& e'_i \longmapsto t_i \otimes e'_i - \hat{e}_i \otimes 1 \\
& f_i \longmapsto -t_i f_i \otimes 1 + t_i \otimes f_i \\
& q^h \longmapsto q^{-h} \otimes q^h \\
\\
\beta_{B^{\text{op}}} : \quad & B^{\text{op}} \longrightarrow B^{\text{op}} \otimes U' \\
& e'_i \longmapsto -t_i^{-1} \otimes e'_i + e'_i \otimes 1 \\
& f_i \longmapsto -t_i^{-1} \otimes f'_i + f_i \otimes 1 \\
& q^h \longmapsto q^h \otimes q^{-h} \\
\\
\gamma : \quad & U' \longrightarrow B \otimes B^{\text{op}} \\
& e'_i \longmapsto t_i e'_i \otimes 1 - t_i \otimes e'_i \\
& f'_i \longmapsto t_i f_i \otimes 1 - t_i \otimes f_i \\
& q^h \longmapsto q^h \otimes q^{-h} \\
\\
\delta : \quad & \hat{U} \longrightarrow B^{\text{op}} \otimes B \\
& \hat{e}_i \longmapsto t_i^{-1} \otimes t_i e'_i - e'_i \otimes 1 \\
& f_i \longmapsto 1 \otimes f_i - f_i t_i \otimes t_i^{-1} \\
& q^h \longmapsto q^{-h} \otimes q^h \\
\\
S_{B^{\text{op}}} := \text{Id}_B : \quad & B^{\text{op}} \longrightarrow (B)^{\text{op}} \\
\\
S_B := \theta_B : \quad & B \longrightarrow (B^{\text{op}})^{\text{op}} = B
\end{aligned}$$

With the help of [G], we can now identify $H_l(B)$ and $H_r(B)$.

Corollary 6 *The morphisms γ and δ define Hopf algebras isomorphisms between U' (resp. $\hat{U}$) and $H_l(B)$ (resp. $H_r(B)$).*

We are now going to extend the scalars to F .

4.4.2 Hopf Galois systems and Kashiwara algebras over F

The quadruple $(B, B^{\text{op}}, U', \hat{U})$ is a complete autonomous Hopf-Galois system. So by [G], we see that $(B_F, B_F^{\text{op}}, U'_F, \hat{U}_F)$ is also a complete autonomous Hopf-Galois system. It follows that $H_r(B_F) \cong \hat{U}_F$ and $H_l(B_F) \cong U'_F$. Therefore, with the help of Corollary 5, we can state the following theorem.

Theorem 5 *Kashiwara algebras equipped with their natural structures of $U_{\hbar}(\mathfrak{g})$ -comodules are $U_{\hbar}(\mathfrak{g})$ -Galois extensions of $\mathbb{Q}((\hbar))$.*

5 Classical limit of B

If we consider the limit when $\hbar \mapsto 0$, then $\hat{e}_i$ and f_j commute (resp. e'_i and f'_j) and the only relations between generators are Serre relations. The classical limit of $\hat{U}$ and the classical limit of U' coincide both with the enveloping algebra $U(\mathfrak{a})$ ($\mathfrak{a}$ has been introduced in the first section). The classical limit B_{cl} of B is the algebra generated by e'_i, f_j ($i, j \in I$) and $h \in \mathfrak{t}$ submitted to relations:

$$[h, e'_i] = \langle \alpha_i, h \rangle e'_i \quad (42)$$

$$[h, f_j] = -\langle \alpha_i, h \rangle f_j \quad (43)$$

$$e'_i f_j = f_j e'_i + \delta_{i,j} \quad (44)$$

together with Serre relations. To the limit when $\hbar \mapsto 0$, we see that the classical limit of Kashiwara algebras B_{cl} are $(U(\mathfrak{a}), U(\mathfrak{a}))$ -quantum torsors over $\mathbb{Q}$. In fact, it is easy to see that classical limit of Kashiwara algebras are examples of Sridharan algebras.

Proposition 5 *The classical limit of Kashiwara algebras B_{cl} are Sridharan algebras for the Lie algebra $\mathfrak{a}$ defined above. The cocycle $c : \mathfrak{a} \wedge \mathfrak{a} \longrightarrow \mathfrak{a}$ is defined by $c(e_i, f_j) = \delta_{i,j}$, $c(e_i, e_j) = c(f_i, f_j) = c(h, e_i) = c(h, f_i) = 0$ and cocycle relations.*

6 Open problems

We finish this article by asking two questions.

6.0.3 Question 1.

If T is a B -Galois extension of k and if the antipode S_B of B is bijective, does this imply θ_T bijective ?

6.0.4 Question 2 (Completion problem).

If (A, B, T, Z) is a Hopf-Galois system in the sense of Bichon, is it possible to find a (B, A) -bicomodule structure on Z and a map $S_T : T \longrightarrow Z$ such that (A, B, T, Z) is a *complete* Hopf-Galois system in the sense we gave in [G] ?

6.1 Acknowledgment

I am very grateful to B. Enriquez for his help and support. I am also grateful to J. Bichon and P. Schauenburg for remarks. I would also like to thank W. Soergel for the hospitality of the Freiburg University.

References

- [B] J. Bichon, Hopf-Galois systems, To appear in Journal of Algebra.
- [G] C. Grunspan, Quantum Torsors, Accepted for publication in Journal of Pure and Applied Algebra.
- [K] M. Kashiwara, On crystal bases of the q -analogue of universal enveloping algebras, Duke Math. J., **63** (1991), 465–516.
- [Sc1] P. Schauenburg, Hopf bi-Galois extensions, Communications in Algebra, **24** (1996), n°12, 3797–3825.
- [Sc2] P. Schauenburg, Quantum torsors and Hopf-Galois objects, preprint, 2002.
- [Sc3] P. Schauenburg, Quantum torsors with fewer axioms, preprint, 2002.
- [Sr] R. Sridharan, Filtered algebras and representations of Lie algebras, Trans. Am. Math. Soc. **100** (1961), 530–550